

Amendments to the Specification

Page 8, line 25 to page 9, line 20

B¹

In Phase I, 60% purified water by weight was used, as it allowed complete blending of all other ingredients while maintaining an ideal consistency. 25.7% aloe vera gel was used, as it provided an acceptable level of anti-inflammatory effect and stimulation of the healing process without a greasy or oily feeling. 8% ~~polyvinol~~ [polyvinyl alcohol] crystals was utilized to increase the exfoliative effect of the product without leaving a sticky feeling after drying. 5% glycerin was used to obtain the correct humectant action. Increased amounts of glycerin tends to impair the exfoliant activity. 0.2% seaweed extract provides improved exfoliation while stabilizing the correct viscosity. 0.2% triclosan was used, as it is the maximum amount allowed by the United States Food and Drug Administration and is needed to achieve the antibacterial action required. A lower dosage would decrease the effectiveness. 0.2% Vitamin B5 (dexapanthanol or pantothenic acid) was used to promote the healing process; an the amount of pantothenic acid is kept low because it can produce local irritation. 0.2% sage extract was used to additionally stimulate the healing process and produce an anti-inflammatory effect, as well as adding to the antibacterial activity. Higher percentage of sage extract may be locally irritating, while a lower amount of sage extract can detract from the healing process and the antibacterial activity. 0.2% chamomile extract was used to supplement the anti-inflammatory effect and speed the healing process and 0.2% rose hips extract was utilized to produce a significant degree of antioxidant activity. Both of these ingredients may cause localized skin irritation unless concentrations are kept relatively low. 0.1% methylparaben was used to increase the range and efficiency of the antibacterial component of the product, as well as stabilizing the product to increase the shelf

B1
Critic

life. Too little methylparaben would result in a decreased shelf life and narrower range of antibacterial activity, while an excess of methylparaben could induce some toxicity factors.

Page 11, lines 20-22

B2
N.E.

The range of ~~polyvinol~~ [polyvinyl alcohol] crystals used allows for maximum exfoliation without producing excess stickiness or skin irritation.
